

---

## UNITED KINGDOM USER PACKAGE LEAFLET OF CERAZETTE

---

RA 0390 GB P8 (ref 5.0)

**Cerazette®**  
**75 microgram film-coated tablet**  
**desogestrel**

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- This leaflet will provide information about the benefits and risks of Cerazette. It will also advise you how to take Cerazette properly and when to tell your doctor about health-related conditions. If you have any further questions, please ask your doctor, pharmacist or Family Planning Nurse.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, pharmacist or Family Planning Nurse.

**In this leaflet:**

- 1. What Cerazette is and what it is used for**
- 2. Before you take Cerazette**  
Do not take...  
Take special care...  
Pregnancy and breast-feeding
- 3. How to take Cerazette**  
Starting your first pack  
    Changing from other methods  
    After a baby  
If you forget to take Cerazette
- 4. Possible side effects**
- 5. How to store Cerazette**
- 6. Further Information**

### **1. WHAT CERAZETTE IS AND WHAT IT IS USED FOR**

- Cerazette is used to prevent pregnancy.
- There are 2 main kinds of hormone contraceptive.
  - The combined pill, "**The Pill**", which contains 2 types of female sex hormone an oestrogen and a progestogen
  - The progestogen-only pill, **POP or mini-pill**, which doesn't contain an oestrogen.
- Cerazette is a progestogen-only-pill (POP), or a **mini-pill**.
- Cerazette contains a small amount of one type of female sex hormone, the progestogen **desogestrel**.
- Most POPs or minipills work primarily by preventing the sperm cells from entering the womb but they do not always prevent the egg cell from ripening, which is the main way that combined pills work.

- Cerazette is different from other mini-pills in having a dose that in most cases prevents the egg cell from ripening. As a result, Cerazette is a highly effective contraceptive.
- In contrast to the combined pill, Cerazette can be used by women who do not tolerate oestrogens and by women who are breast feeding.
- A disadvantage is that vaginal bleeding may occur at irregular intervals during the use of Cerazette. On the other hand you may not have any bleeding at all.

## 2. BEFORE YOU TAKE CERAZETTE

Cerazette, like other hormonal contraceptives, does not protect against HIV infection (AIDS) or any other sexually transmitted disease.

### 2.1 Do not take Cerazette

- if you are **allergic** to any of the ingredients of Cerazette.
- if you have a thrombosis. Thrombosis is the formation of a **blood clot** in a blood vessel (e.g. of the legs (deep venous thrombosis) or the lungs (pulmonary embolism)).
- if you have or have had **jaundice** (yellowing of the skin) or severe liver disease and your liver is still not working normally.
- if you have or if you are suspected of having a **cancer** that grows under the influence of sex-steroids, such as certain types of breast cancer.
- if you have any unexplained **vaginal bleeding**.

If any of these conditions apply to you, tell your doctor before you start to use Cerazette. Your doctor may advise you to use a non-hormonal method of birth control.

If any of these conditions appear for the first time while using Cerazette, consult your doctor immediately.

### 2.2 Take special care with Cerazette

Before you start Cerazette tell your doctor or Family Planning Nurse, if

- you have ever had **breast cancer**.
- you have **liver cancer**, since a possible effect of Cerazette cannot be excluded.
- you have ever had a **thrombosis**.
- you have **diabetes**.
- you suffer from **epilepsy** (see section 'Taking other medicines').
- you have **tuberculosis** (see section 'Taking other medicines').
- you have high **blood pressure**.
- you have or have had **chloasma** (yellowish-brown pigmentation patches on the skin, particularly of the face); if so, avoid too much exposure to the sun or ultraviolet radiation.

When Cerazette is used in the presence of any of these conditions, you may need to be kept under close observation. Your doctor can explain what to do.

### 2.3 Breast cancer

- It is important to regularly check your breasts and you should contact your doctor as soon as possible if you feel a lump in your breasts.
- Breast cancer has been found slightly more often in women who take the Pill than in women of the same age who do not take the Pill. If women stop taking the Pill, this reduces the risk, so that 10 years after stopping the Pill, the risk is the same as for women who have never taken the Pill.

Breast cancer is rare under 40 years of age but the risk increases as the woman gets older. Therefore, the extra number of breast cancers diagnosed is higher if a woman continues to take the Pill when she is older. How long she takes the Pill is less important.

- In every 10 000 women who take the Pill for up to 5 years but stop taking it by the age of 20, there would be less than 1 extra case of breast cancer found up to 10 years after stopping, in addition to the 4 cases normally diagnosed in this age group.
- In 10 000 women who take the Pill for up to 5 years but stop taking it by the age of 30, there would be 5 extra cases in addition to the 44 cases normally diagnosed.
- In 10 000 women who take the Pill for up to 5 years but stop taking it by the age of 40, there would be 20 extra cases in addition to the 160 cases normally diagnosed.

The risk of breast cancer in users of progestogen-only pills like Cerazette is believed to be similar to that in women who use the Pill, but the evidence is less conclusive.

Breast cancers found in women who take the Pill, seem less likely to have spread than breast cancers found in women who do not take the Pill.

It is not certain whether the Pill causes the increased risk of breast cancer. It may be that the women were examined more often, so that the breast cancer is noticed earlier.

#### **2.4 Thrombosis**

See your doctor immediately if you notice possible signs of a thrombosis (see also ‘Regular check-ups’).

Thrombosis is the formation of a **blood clot**, which may block a blood vessel. A thrombosis sometimes occurs in the deep veins of the legs (deep venous thrombosis). If this clot breaks away from the veins where it is formed, it may reach and block the arteries of the lungs, causing a so-called “pulmonary embolism”. A pulmonary embolism can cause chest pain, breathlessness, collapse or even death.

- Deep venous thrombosis is a rare occurrence. It can develop whether or not you are taking the Pill. It can also happen if you become pregnant.

The risk is higher in Pill-users than in non-users. The risk with progestogen-only pills like Cerazette is believed to be lower than in users of Pills that also contain oestrogens (combined Pills).

#### **2.5 Taking other medicines**

Please tell your doctor, pharmacist, or Family Planning Nurse if you are taking or have recently taken any other medicines or herbal products, including medicines obtained without a prescription.

Some medicines may stop Cerazette from working properly. These include medicines used for the treatment of

- epilepsy (e.g. primidone, phenytoin, carbamazepine, oxcarbazepine, felbamate and phenobarbital)
- tuberculosis (e.g. rifampicin)
- HIV infections (e.g. ritonavir), or other infectious diseases (e.g. griseofulvin)
- stomach upset (medical charcoal)

- depressive moods (the herbal remedy St. John's Wort).

Your doctor can tell you if you need to take additional contraceptive precautions and if so, for how long.

Cerazette may also interfere with how certain medicines work, causing either an increase in effect (e.g. medicines containing cyclosporine) or a decrease in effect.

## **2.6 Pregnancy and breast-feeding**

### Pregnancy

Do not use Cerazette if you are pregnant, or think you may be pregnant.

### Breast-feeding

Cerazette may be used while you are breast-feeding. Cerazette does not influence the production or the quality of breast milk. However, a small amount of the active substance of Cerazette passes over into the milk.

The health of children who were breast-fed for 7 months while their mothers were using Cerazette has been studied up until they were 2½ years of age. No effects on the growth and development of the children were observed.

If you are breast feeding and want to use Cerazette, please contact your doctor.

## **2.7 Driving and using machines**

Cerazette has no known effect on the ability to drive or use machines

## **2.8 Important information about some of the ingredients of Cerazette**

Cerazette contains lactose (milk sugar). Please contact your doctor before taking Cerazette if you have been told by your doctor that you are intolerant to some sugars.

## **2.9 Use in adolescents**

No clinical data on efficacy and safety are available in adolescents below 18 years.

## **2.10 Regular check-ups**

When you are using Cerazette, your doctor will tell you to return for regular check-ups. In general, the frequency and nature of these check-ups will depend on your personal situation.

### *Contact your doctor as soon as possible if:*

- you notice possible signs of a **blood clot** e.g. severe pain or swelling in either of your legs; unexplained pains in the chest, breathlessness, an unusual cough, especially when you cough up blood;
- you have a sudden, severe stomach ache or **jaundice** (you may notice yellowing of the skin, the whites of the eyes, or dark urine, possibly a sign of **liver problems**);
- you feel a lump in your **breast**;
- you have a sudden or severe pain in the lower abdomen or stomach area (possibly a sign of an **ectopic pregnancy** - a pregnancy outside the womb);
- you are to be immobilised or are to have surgery (consult your doctor at least four weeks in advance);
- you have unusual, heavy **vaginal bleeding**;
- you suspect that you are **pregnant**.

### 3. HOW TO TAKE CERAZETTE

#### 3.1 When and how to take the tablets?

Each strip of Cerazette contains 28 tablets – 4 weeks supply.

- **Take your tablet each day at about the same time. Swallow the tablet whole, with water.**
- Arrows are printed on the front of the strip, between the tablets. The days of the week are printed on the back of the strip. Each day corresponds with one tablet.
- Every time you start a new strip of Cerazette, take a tablet from the top row. Don't start with just any tablet. For example if you start on a Wednesday, you must take the tablet from the top row marked (on the back) with WED.
- Continue to take one tablet a every day until the pack is empty, always following the direction indicated by the arrows. By looking at the back of your pack you can easily check if you have already taken your tablet on a particular day.
- You may have some vaginal bleeding during the use of Cerazette, (See Section 4 Side Effects) but you must continue to take your tablets as normal.
- When a strip is empty, you must start with a new strip of Cerazette on the next day - without interruption and without waiting for a bleed.

#### 3.2 Starting your first pack of Cerazette

- **If you are not using hormonal contraception at present (or in the past month)**

Wait for your period to begin. On the first day of your period take the first Cerazette tablet.

Additional contraceptive precautions are not necessary. If you take your first tablet on days 2-5 of your period use an additional barrier method of contraception for the first 7 days of tablet-taking.

- **When you change from a combined pill (COC), vaginal ring, or transdermal patch**

**If you have** a tablet-, ring- or patch-free break

- You can also start at the latest the day following the tablet-, ring- or patch-free break, or placebo tablet interval, of your present contraceptive.
- **If you follow these instructions, make sure you use an additional barrier method of contraception for the first 7 days of tablet-taking.**

**If you don't have** a tablet-, ring- or patch-free break

- Start taking Cerazette on the day after you take the last tablet from the present Pill pack, or on the day of removal of your vaginal ring or patch (this means no tablet-, ring- or patch-free break).
- If your present Pill pack also contains inactive tablets you can start Cerazette on the day after taking the last active tablet (if you are not sure which this is, ask your doctor or pharmacist).
- **If you follow these instructions, additional contraceptive precautions are not necessary.**

- **When changing from another progestogen-only pill (mini-pill)**

Switch on any day from another mini pill. Additional contraceptive precautions are not necessary.

- **When changing from an injection or implant or a hormonal IUS**

Start using Cerazette when your next injection is due or on the day that your implant or your IUS is removed. Additional contraceptive precautions are not necessary.

- **After you have a baby**

You can start Cerazette between 21 to 28 days after the birth of your baby.

If you start later, make sure that during the first cycle you use an additional barrier method of contraception for the first 7 days of tablet-taking. However, if you have already had sex, check that you are not pregnant before starting Cerazette. Information for breast-feeding women can be found in section 2 'Before you take Cerazette' in the paragraph 'Pregnancy and breast-feeding'. Your doctor can also advise you.

- **After a miscarriage or an abortion**

Your doctor will advise you.

### **3.3 If you forget to take Cerazette**

- If you are **less than 12 hours** late:
  - Take the missed tablet as soon as you remember and take the next one at the usual time. Cerazette will still protect you from pregnancy.
- If you are **more than 12 hours** late:
  - If you are more than 12 hours late in taking any tablet, **you may not be completely protected against pregnancy**. The more consecutive tablets you have missed, the higher the risk that you might fall pregnant.
  - Take a tablet as soon as you remember and take the next one at the usual time. This may mean taking two in one day. This is not harmful. (If you have forgotten more than one tablet you don't need to take the earlier missed ones). Continue to take your tablets as usual but you must also use an extra method, such as a condom, for the **next 7 days**.
  - If you are more than 12 hours late taking your tablet and have had sex it is safe to use **emergency contraception**; please consult your pharmacist or doctor.
  - If you missed one or more tablets in the **very first week** of tablet-intake and had intercourse in the week before missing the tablets, you may fall pregnant. Ask your doctor for advice.

### **3.4 If you vomit or use medical charcoal**

If you vomit, or use medical charcoal within 3 - 4 hours after taking your Cerazette tablet or have severe diarrhoea, the active ingredient may not have been completely absorbed. Follow the advice for forgotten tablets in the section above.

### **3.5 If too many Cerazette tablets are taken (overdose)**

There have been no reports of serious harmful effects from taking too many Cerazette tablets at one time. Symptoms that may occur are nausea, vomiting and in young girls, slight vaginal bleeding. For more information ask your doctor for advice.

### **3.6 If you stop taking Cerazette**

You can stop taking Cerazette whenever you want. From the day you stop you are no longer protected against pregnancy.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

#### 4. POSSIBLE SIDE EFFECTS

Like all medicines, Cerazette can cause side effects, although not everybody gets them. Tell your doctor if you notice any unwanted effect, especially if severe or persistent.

**Serious side effects** associated with the use of Cerazette are described in section 2 'Before you take Cerazette'. Please read this section for additional information on 'Breast cancer' and 'Thrombosis' and consult your doctor at once where appropriate.

**Vaginal bleeding may occur at irregular intervals** while using Cerazette. This may be just slight staining which may not even require a pad, or heavier bleeding, which looks rather like a scanty period. You may need to use tampons or sanitary towels. You may also not have any bleeding at all. Irregular bleeding is not a sign that Cerazette is not working. In general, you need not take any action; just continue to take Cerazette. **If bleeding is heavy or prolonged you should consult your doctor.**

#### How often are other possible side effects seen?

**Common** (affecting less than 1 in 10 women): mood changes, depressed mood, decreased sexual drive (libido), headache, nausea, acne, breast pain, irregular or no periods, weight increase.

**Uncommon** (affecting less than 1 in 100 women) infection of the vagina, difficulties in wearing contact lenses, vomiting, hair loss, painful periods, ovarian cysts, tiredness.

**Rare** (affecting less than 1 in 1000 women) skin conditions such as: rash, hives, painful blue-red skin lumps (erythema nodosum)

Apart from these side effects, breast secretion or leakage may occur.

You should see your doctor immediately if you experience symptoms of angioedema, such as (i) swollen face, tongue or pharynx; (ii) difficulty to swallow; or (iii) hives and difficulties to breathe.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, pharmacist or Family Planning Nurse.

#### 5. HOW TO STORE CERAZETTE

Keep out of the reach and sight of children.

Do not use after the expiry date which is stated on the package. The expiry date is the last day of the month stated. Cerazette does not require any special storage conditions. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### 6. FURTHER INFORMATION

##### What Cerazette contains

- The active substance is: desogestrel (75 microgram)
- The other ingredients are: colloidal anhydrous silica; all-rac- $\alpha$ -tocopherol; maize starch; povidone; stearic acid; hypromellose; macrogol 400; talc; titanium dioxide (E171); lactose monohydrate (see also 'Important information about some of the ingredients of Cerazette' in section 2).

**What Cerazette looks like and contents of the pack**

Each strip of Cerazette contains 28 white round tablets. The tablets are marked KV above 2 on one side and ORGANON\* on the other. Cerazette comes in a pack of 3 strips. Each strip is sealed in a foil sachet.

**Marketing Authorisation Holder**

*Organon Laboratories Ltd, Science Park, Milton Road, Cambridge CB4 0FL, UK*

**Manufacturer:**

N.V. Organon, P.O. Box 20, 5340 BH Oss, The Netherlands, or  
Organon (Ireland) Ltd., Drynam Road, P.O. Box 2857, Swords, Co. Dublin, Ireland

**This leaflet was approved in January 2012**

In correspondence please quote packing number.

To listen to or request a copy of this leaflet in **Braille, large print or audio** please call, free of charge: 0800 198 5000 (UK Only)

Please be ready to give the following information:

Product name: Cerazette                      Reference Number: PL 00065/0159

This is a service provided by the Royal National Institute of Blind people.