

**Bayer plc**

Bayer House, Strawberry Hill, Newbury, Berkshire, RG14 1JA

Telephone: +44 (0)1635 563 000

Fax: +44 (0)1635 563 393

WWW: <http://www.bayer.co.uk>



Due to regulatory changes, the content of the following Patient Information Leaflet may vary from the one found in your medicine pack. Please compare the 'Leaflet prepared/revised date' towards the end of the leaflet to establish if there have been any changes.

If you have any doubts or queries about your medication, please contact your doctor or pharmacist.

**Package Leaflet: Information for the user**  
**Levitra 10 mg film-coated tablets**  
Vardenafil

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- *Keep this leaflet. You may need to read it again.*
- *If you have any further questions, ask your doctor or pharmacist.*
- *This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.*
- *If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.*

**What is in this leaflet:**

1. *What Levitra is and what it is used for*
2. *What you need to know before you take Levitra*
3. *How to take Levitra*
4. *Possible side effects*
5. *How to store Levitra*
6. *Contents of the pack and other information*

**1. What Levitra is and what it is used for**

*Levitra contains vardenafil, a member of a class of medicines called phosphodiesterase type 5 inhibitors. They are used for the treatment of erectile dysfunction in adult men, a condition which implies difficulties in getting or keeping an erection.*

*At least one in ten men has trouble getting or keeping an erection at some time. There may be physical or psychological causes, or a mixture of both. Whatever the cause is, due to muscle and blood vessel changes not enough blood stays in the penis to make it hard and keep it hard.*

*Levitra will only work when you are sexually stimulated. It reduces the action of the natural chemical in your body which makes erections go away. Levitra allows an erection to last long enough for you to satisfactorily complete sexual activity.*

## **2. What you need to know before you take Levitra**

### **Do not take Levitra**

- If you are allergic to vardenafil or any of the other ingredients of this medicine (listed in section 6). Signs of an allergic reaction include a rash, itching, swollen face or lips and shortness of breath.
- If you are taking medicines containing nitrates, such as glycerol trinitrate for angina, or nitric oxide donors, such as amyl nitrite. Taking these medicines with Levitra could seriously affect your blood pressure.
- If you are taking ritonavir or indinavir, medicines used to treat human immunodeficiency virus (HIV) infections.
- If you are over 75 years of age and are taking ketoconazole or itraconazole, anti-fungal medicines.
- If you have a severe heart or liver problem.
- If you are having kidney dialysis.
- If you have recently had a stroke or heart attack.
- If you have or have had low blood pressure.
- If your family has a history of degenerative eye diseases (such as retinitis pigmentosa).
- If you have ever had a condition involving [loss of vision](#) due to damage to the [optic nerve](#) from insufficient blood supply known as non-arteritic ischemic optic neuropathy (NAION).

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Levitra.

### **Take special care with Levitra**

- If you have heart trouble. It may be risky for you to have sex.
- If you suffer from irregular heart beat (cardiac arrhythmia) or inherited heart diseases affecting your electrocardiogram.
- If you have a physical condition affecting the shape of the penis. This includes conditions called angulation, Peyronie's disease and cavernosal fibrosis.
- If you have an illness that can cause erections which won't go away (priapism). These include sickle cell disease, multiple myeloma and leukaemia.
- If you have stomach ulcers (also called gastric or peptic ulcers).
- If you have a bleeding disorder (such as haemophilia).
- If you are using any other treatments for erection difficulties, including Levitra orodispersible tablets. (see section Other medicines and Levitra).
- If you experience sudden decrease or loss of vision, stop taking Levitra and contact your doctor immediately.

### **Children and adolescents**

Levitra is not intended for use by children or adolescents under 18.

### **Other medicines and Levitra**

Please tell your doctor or pharmacist if you are using, have recently used or might use any other medicines, including medicines obtained without a prescription.

Some medicines may cause problems, especially these:

- Nitrates, medicines for angina, or nitric oxide donors, such as amyl nitrite. Taking these medicines with Levitra could seriously affect your blood pressure. Talk to a doctor without taking Levitra.
- Medicine for the treatment of arrhythmias, such as quinidine, procainamide, amiodarone or sotalol.
- Ritonavir or indinavir, medicines for HIV. Talk to a doctor without taking Levitra.
- Ketoconazole or itraconazole, anti-fungal medicines
- Erythromycin or clarithromycin, macrolide antibiotics
- Alpha-blockers, a type of medicine used to treat high blood pressure and enlargement of the prostate (as benign prostatic hyperplasia).

Do not use Levitra film-coated tablets combined with any other treatment for erectile dysfunction, including Levitra orodispersible tablets.

### **Taking Levitra with food and drink**

- You can take Levitra with or without food – but preferably not after a heavy or high-fat meal as this may delay the effect.
- Don't drink grapefruit juice when you use Levitra. It can interfere with the usual effect of the medicine.
- Alcoholic drink can make erection difficulties worse.

### **Pregnancy and breast-feeding**

Levitra is not for use by women.

### **Driving and using machines**

Levitra might make some people feel dizzy or affect their vision. If you feel dizzy, or if your vision is affected after taking Levitra don't drive or operate any tools or machines.

### **3. How to take Levitra**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. The recommended dose is 10 mg.

Take a Levitra tablet about 25 to 60 minutes before sexual activity. With sexual stimulation you may achieve an erection anywhere from 25 minutes up to four to five hours after taking Levitra.

- Swallow one tablet with a glass of water

**Do not take Levitra film-coated tablets** with any other forms of Levitra.

**Do not take Levitra** more than once a day.

Tell your doctor if you think Levitra is too strong or too weak. He or she may suggest a switch to an alternative Levitra formulation with a different dose, depending on how well it works for you.

#### **If you take more Levitra than you should**

Men who take too much Levitra may experience more side effects or may get severe back pain. If you take more Levitra than you should, tell your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the effects are mild or moderate.

Partial, sudden, temporary or permanent decrease or loss of vision in one or both eyes has been experienced by patients. Stop taking Levitra and contact your doctor immediately.

Sudden decrease or loss of hearing has been reported.

The chance of having a side effect is described by the following categories:

#### **Very common:**

may affect more than 1 in 10 users

- Headache

#### **Common:**

may affect up to 1 in 10 users

- Dizziness
- Flushing
- Blocked or runny nose
- Indigestion

#### **Uncommon:**

may affect 1 in 100 users

- Swelling of skin and mucous tissue including swollen face, lips or throat
- Sleep disorder
- Numbness and impaired perception of touch
- Sleepiness

- *Effects on vision; redness of the eye, effects on colour vision, eye pain and discomfort, light sensitivity*
- *Ringing in the ears; vertigo*
- *Fast heart beat or pounding heart*
- *Breathlessness*
- *Stuffy nose*
- *Acid reflux, gastritis, abdominal pain, diarrhoea, vomiting; feeling sick (nausea), dry mouth*
- *Raised levels of liver enzymes in your blood*
- *Rash, reddened skin*
- *Back or muscle pain; increase in blood of a muscle enzyme (creatine phosphokinase), muscle stiffness*
- *Prolonged erections*
- *Malaise*

**Rare:**

*may affect 1 in 1,000 users*

- *Inflammation of the eyes (conjunctivitis)*
- *Allergic reaction*
- *Anxiety*
- *Fainting*
- *Amnesia*
- *Seizure*
- *Increase pressure in the eye (glaucoma), lacrimation increased*
- *Effects on the heart (such as heart attack, altered heart beat or angina)*
- *High or low blood pressure*
- *Nose bleed*
- *Effect on results of blood tests to check liver function*
- *Sensitivity of the skin to sun light*
- *Painful erections*
- *Chest pain*

*If any of the side effects gets serious or if you notice any side effects not listed in this leaflet, please tell your doctor.*

**5. How to store Levitra**

*Keep this medicine out of the sight and reach of children.*

*Do not use this medicine after the expiry date which is stated on the carton after “EXP”. The expiry date refers to the last day of that month.*

*Do not throw away any medicines of via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.*

**6. Contents of the pack and other information**

**What Levitra contains**

- *The active substance is vardenafil. Each tablet contains 10 mg of vardenafil (as hydrochloride ).*
- *The other ingredients of the tablets are:*  
*Tablet core: crospovidone, magnesium stearate, microcrystalline cellulose, colloidal anhydrous silica.*  
*Film coat: macrogol 400, hypromellose, titanium dioxide (E171), ferric oxide yellow (E172), ferric oxide red (E172).*

**What Levitra looks like and contents of the pack**

*Levitra 10 mg film-coated tablets are orange with the BAYER cross on one side and the strength (10) on the other side. The tablets are provided in blister packs containing 2, 4, 8, 12 or 20 tablets. Not all pack sizes may be marketed.*

***Marketing Authorisation Holder and Manufacturer***

*Marketing Authorisation Holder*

*Bayer Pharma AG*

*13342 Berlin*

*Germany*

*Manufacturer*

*Bayer Pharma AG*

*51368 Leverkusen*

*Germany*

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Bayer S.A./N.V. Tél/Tel:  
+32-(0)2-535 63 11

**България**

Байер България ЕООД  
+359 02 81 401 01

**Česká republika**

GlaxoSmithKline s.r.o.  
Tel: +420 2 22001111

**Danmark**

Bayer A/S  
Tlf: +45-45-23 50 00

**Deutschland**

Jenapharm GmbH & Co. KG  
Tel: +49-(0)3641-64 8888

**Eesti**

GlaxoSmithKline Eesti OÜ  
Tel: +372 6676900  
Estonia@GSK.com

**Ελλάδα**

Bayer Ελλάς ABEE Τηλ:  
: +30 210 618 75 00

**España**

Bayer Hispania S.L.  
Tel: +34-93-495 65 00

**France**

Bayer Santé Tél:  
+33-(0)3-28 16 34 00

**Ireland**

Bayer Limited  
Tel: +353 1 299 93 13

**Ísland**

Icepharma hf. Sími:  
+354-540 8000

**Italia**

Bayer S.p.A.  
Tel: +39-02-397 81

**Κύπρος**

NOVAGEM Limited  
Τηλ.: +357 22 48 38 58

**Latvija**

GlaxoSmithKline Latvia SIA  
Tel: +371-67312687

**Lietuva**

UAB "GlaxoSmithKline Lietuva"  
Tel: +370 5 264 90 00

**Luxembourg/Luxemburg**

Bayer S.A./N.V. Tél/Tel:  
+32-(0)2-535 63 11

**Magyarország**

Bayer Hungária Kft.  
Tel.: +36-1-487-4100

**Malta**

Alfred Gera and Sons Ltd. Tel:  
+356-21 44 62 05

**Nederland**

Bayer B.V. Tel:  
+31-(0)297-28 06 66

**Norge**

Bayer AS Tlf:  
+47 24 11 18 00

**Österreich**

Bayer Austria Ges. m. b. H.  
Tel: +43-(0)1-711 46-0

**Polska**

Bayer Sp. z o.o.  
Tel.: +48-22-572 35 00

**Portugal**

Bayer Portugal S.A.  
Tel: +351-21-416 42 00

**România**

GlaxoSmithKline (GSK) S.R.L. Tel:  
+4021 3028 208

**Slovenija**

Bayer d. o. o.  
Tel: +386 1 58 14 400

**Slovenská republika**

GlaxoSmithKline Slovakia s.r.o. Tel:  
+421 (0)2 48261111

**Suomi/Finland**

Bayer Oy  
Puh/Tel: +358-20 785 21

**Sverige**

Bayer AB  
Tel: +46 (0)8 580 223 00

**United Kingdom**

Bayer plc  
Tel: +44-(0)1635-563000

*This leaflet was last revised in 03/2012*

Detailed information on this medicinal product is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

